

Contribution ID : 32

Type : Oral

Exotic interactions mediated by a non-Hermitian photonic bath

mercoledì 13 ottobre 2021 15:45 (25)

We study the exotic interaction between emitters mediated by the photonic modes of a lossy photonic lattice described by a non-Hermitian Hamiltonian, where structured losses can seed exotic emission properties. Photons can mediate dissipative, fully non-reciprocal, interactions between emitters with range critically dependent on the loss rate. At the bare-lattice exceptional point, the effective couplings are exactly nearestneighbour, implementing a dissipative, fully non-reciprocal, Hatano-Nelson model. Counter-intuitively, this occurs irrespective of the lattice boundary conditions. Thus photons can mediate an effective emitters' Hamiltonian which is translationally-invariant despite the fact that the field is not. We interpret these effects in terms of metastable atom-photon dressed states, which can be exactly localized on only two lattice cells or extended across the entire lattice.

Primary author(s): ROCCATI, Federico (Università degli Studi di Palermo)

Co-author(s) : LORENZO, Salvatore (Università degli Studi di Palermo); CALAJÒ, Giuseppe (ICFO); PALMA, Massimo (Università degli Studi di Palermo); CAROLLO, Angelo (Università degli Studi di Palermo); CICCARELLO, Francesco (Università degli Studi di Palermo)

Presenter(s) : ROCCATI, Federico (Università degli Studi di Palermo)

Session Classification : Session 6