



Contribution ID : 5

Type : not specified

Terahertz saturable absorption from relativistic high-temperature thermodynamics in black phosphorus

Tuesday, 20 February 2024 15:00 (40)

Black phosphorus is a unique two-dimensional (2D) material (Figure 1) with a tunable infrared band gap and anisotropic conduction properties [1]. Black phosphorus also displays the occurrence of a pressure-induced topological Lifshitz transition turning the material from a narrow gap semiconductor to a massless Dirac metal due to a nonavoided band crossing [2]. We investigate the ambient pressure nonlinear terahertz (THz) electrodynamic properties of black phosphorus along the more conducting armchair direction and found that its THz saturable-absorption properties can be understood within a thermodynamic model by assuming a fast thermalization of the electron bath [3]. While black phosphorus does not display the presence of massless fermions at ambient pressure and temperature the material's anomalous THz nonlinear properties can be accounted for by a relativistic massive Dirac dispersion, provided that the Fermi temperature is low enough. This suggests that an optimal tuning of the Fermi level could be a strategy to engineer a strong THz nonlinear response in other massive Dirac materials, such as transition-metal dichalcogenides or high-temperature superconductors

Primary author(s) : ADHLAKHA, Nidhi (Elettra-Sincrotrone Trieste); EBRAHIMPOUR, Zeinab; DI PIETRO, Paola; SCHMIDT, Johannes; PICCIRILLI, Federica; FAUSTI, Daniele; MONTANARO, Angela; CAPPELLUTI, Emmanuele; LUPI, Stefano; PERUCCHI, Andrea

Presenter(s) : PERUCCHI, Andrea

Session Classification : VI session